

Product Information

Anti-Histone Deacetylase 6 (HDAC6)

Developed in Rabbit
IgG Fraction of Antiserum

Product Number **H 2287**

Product Description

Anti-Histone Deacetylase 6 (HDAC6), is developed in rabbit using a synthetic peptide corresponding to amino acid residues 1199-1213 of human HDAC6, conjugated to KLH as immunogen. The corresponding sequence in mouse differs by one amino acid. Whole antiserum is fractionated and then further purified by ion-exchange chromatography to provide the IgG fraction of antiserum that is essentially free of other rabbit serum proteins.

Anti-Histone Deacetylase 6 (HDAC6) recognizes HDAC6 (approx. 134 kDa). Applications include immunoblotting and immunoprecipitation. Detection of HDAC6 by immunoblotting is specifically inhibited with the immunizing peptide. Additional non-specific bands of approx. 50 kDa may be detected in various extract preparations.

Regulation of gene expression is mediated by several mechanisms. Among them are DNA methylation, ATP-dependent chromatin remodeling, and posttranslational modifications of histones, such as the dynamic acetylation and deacetylation of ϵ -amino groups of lysine residues present in the tail of core histones.¹ The enzymes responsible for this reversible acetylation/-deacetylation process are histone acetyltransferases (HATs) and histone deacetylases (HDACs), respectively.² HATs act as transcriptional coactivators and HDACs are part of transcriptional corepressor complexes.³

Mammalian HDACs can be divided into three classes according to sequence homology.⁴ Class I consists of the yeast Rpd3-like proteins HDAC1, HDAC2, HDAC3 and HDAC8. Class II consists of the yeast Hda1-like proteins HDAC4, HDAC5, HDAC6, HDAC7, HDAC9 and HDAC10.⁵ Class III consists of the yeast Sir2-like proteins. Whereas class I HDACs are ubiquitously expressed, most class II HDACs are tissue-specific.² Class II HDACs are larger than those of class I and their catalytic domain is located in the carboxy-terminal half of the protein.

HDAC6 is a unique deacetylase among class II members in that it contains a duplicate of the catalytic domain in its NH₂ terminus as well.² The deacetylase activity of class II HDACs is regulated by subcellular localization.⁴ Endogenous HDAC6 is predominantly cytoplasmic where it associates with microtubules and functions as an α -tubulin deacetylase.⁶ HDAC6 contains a conserved zinc finger motif that is probably involved in the regulation of ubiquitination.⁷

Reagent

Anti-Histone Deacetylase 6 (HDAC6) is supplied as a solution in 0.01 M phosphate buffered saline, pH 7.4, containing 15 mM sodium azide.

Precautions and Disclaimer

Due to the sodium azide content, a material safety data sheet (MSDS) for this product has been sent to the attention of the safety officer of your institution. Consult the MSDS for information regarding hazards and safe handling practices.

Storage/Stability

For continuous use, store at 2-8 °C for up to one month. For prolonged storage, freeze in working aliquots at -20 °C. Repeated freezing and thawing is not recommended. Storage in frost-free freezers is also not recommended. If slight turbidity occurs upon prolonged storage, clarify the solution by centrifugation before use. Working dilutions should be discarded if not used within 12 hours.

Product Profile

For immunoblotting, a minimum working antibody dilution of 1:1,000 is recommended using 293T cells expressing recombinant mouse HDAC6 in a chemiluminescent detection system.

For immunoprecipitation, 5-10 μ g of the antibody immunoprecipitates HDAC6 from a RIPA extract of 500 μ g of 293T cells expressing recombinant mouse HDAC6.

Note: In order to obtain the best results using various techniques and preparations, we recommend determining the optimal working dilutions by titration.

References

1. Wang, A.H., et al., *Mol. Cell. Biol.*, **19**, 7816-7827 (1999).
2. Grozinger, C.M., et al., *Proc. Natl. Acad. Sci. USA*, **96**, 4868-4873 (1999).
3. Fischle, W., et al., *Biochem. Cell Biol.*, **79**, 337-348 (2001).
4. Khochbin, S., et al., *Curr. Opin. Genet. Dev.*, **11**, 162-166 (2001).
5. Fischle, W., et al., *J. Biol. Chem.*, **274**, 11713-11720 (1999).
6. Hubbert, C., et al., *Nature*, **417**, 455-458 (2002).
7. Seigneurin-Berny, D., et al., *Mol. Cell. Biol.*, **21**, 8035-8044 (2001).

ST/KAA 10/02

Sigma brand products are sold through Sigma-Aldrich, Inc.

Sigma-Aldrich, Inc. warrants that its products conform to the information contained in this and other Sigma-Aldrich publications. Purchaser must determine the suitability of the product(s) for their particular use. Additional terms and conditions may apply. Please see reverse side of the invoice or packing slip.