

# Product Information

sigma-aldrich.com

3050 Spruce Street, Saint Louis, MO 63103 USA  
Tel: (800) 521-8956 (314) 771-5765 Fax: (800) 325-5052 (314) 771-5757  
email: techservice@sial.com sigma-aldrich.com

## Anti-Keratinocyte Growth Factor

produced in goat, affinity isolated antibody

Catalog Number **K4760**

Synonyms: Anti-KGF; Anti-FGF-7

### Product Description

Anti-Keratinocyte Growth Factor is produced in goat using as immunogen a purified recombinant human KGF/fibroblast growth factor 7 expressed in *E. coli*. FGF-7 specific IgG was purified by human FGF-7 affinity chromatography.

Anti-Keratinocyte Growth Factor recognizes recombinant human FGF-7 by various immunochemical techniques including neutralization, immunoblotting, immunohistochemistry, and ELISA. Based on ELISA and immunoblotting, this antibody shows less than 20% cross-reactivity with recombinant human FGF-9 and less than 10% cross-reactivity with recombinant human FGF-4.

Fibroblast growth factors (FGFs) are members of a large family of structurally related polypeptides (17-38 kDa) that exert biological activities toward cells of mesenchymal, neuronal, and epithelial origin.<sup>1,2</sup> All members of the FGF superfamily have two conserved cysteine residues and a conserved 120 amino acid core region that contains six identical, interspersed amino acids.<sup>3-5</sup> All FGFs share 30-50 % amino acid sequence identity. FGFs are involved in normal development, wound healing and repair, angiogenesis, and a variety of neurotrophic activities. They are also involved in hematopoiesis as well as in tissue remodeling and maintenance. FGFs are potent physiological regulators of growth and differentiation for a variety of cells of mesodermal, ectodermal, and endodermal origin. They have been implicated in pathological conditions such as tumorigenesis and metastasis. To date, the FGF family consists of 23 members designated FGF-1 through FGF-23.<sup>5</sup>

Four distinct tyrosine kinase FGF receptors (FGFRs) from four separate genes have been identified: FGFR-1 (flg, cek-1), FGFR-2 (bek, cek-3), FGFR-3 (cek-2), and FGFR-4.<sup>6-8</sup> The high affinity cell surface FGF receptors have an extracellular region containing three immunoglobulin-like domains, a transmembrane region, and a cytosolic tyrosine kinase domain activated by

ligand binding. Multiple additional variants (isoforms) arising from alternative splicing have also been reported.<sup>7</sup> Ligand binding specificity, signal transduction, and membrane attachment may be modified by alternative splicings.

Keratinocyte Growth Factor (KGF, FGF-7) was originally isolated from the conditioned medium of a human embryonic lung fibroblast cell line as a mitogen specific for epithelial cells. The transcript for KGF is detected in stromal but not epithelial cells. It has been suggested that KGF is a mesenchymal cell-derived paracrine growth factor that stimulates epithelial cell growth. The N-terminal residues (31 amino acid residues) of KGF are cleaved from the KGF precursor protein (194 amino acid residues) to generate the mature form of KGF (163 amino acid residues).<sup>9</sup> Human KGF shows species cross-reactivity and is active on mouse, monkey, and porcine cells. Mature human and mouse FGF-7 have 96 % amino acid sequence identity, while human and rat FGF-7 have 92 % amino acid sequence identity.<sup>9</sup>

KGF is expressed in a number of stromal fibroblast cell lines but is absent from normal glial cells and epithelial cell lines. KGF released from stromal and dermal fibroblasts stimulates proliferation and differentiation of keratinocytes and other cells in the epithelium. Adult cells known to express FGF-7 include fibroblasts,  $\gamma\delta$  T cells, smooth muscle cells, and ovarian theca cells. In the embryo, KGF is expressed at many stages of development throughout the mesenchyme.

### Reagent

Supplied lyophilized from a 0.2  $\mu$ m filtered solution of phosphate buffered saline with 5% trehalose.

### Preparation Instructions

To one vial of lyophilized powder, add 1 mL of sterile phosphate buffered saline, pH 7.4, to produce a 0.25 mg/mL stock solution of antibody.

### Storage/Stability

Prior to reconstitution, store at  $-20^{\circ}\text{C}$ . Reconstituted product may be stored at  $2-8^{\circ}\text{C}$  for up to one month without detectable loss of activity. For prolonged storage, freeze in working aliquots at  $-20^{\circ}\text{C}$ . Avoid repeated freezing and thawing. Do not store in a frost-free freezer.

### Product Profile

Anti-Keratinocyte Growth Factor has the ability to neutralize the biological activity of recombinant human FGF-7 (KGF) in monkey epithelial 4MBr-5 cells. Recombinant human FGF-7 is added to various concentrations of the antibody for 1 hour at  $22^{\circ}\text{C}$  in a 96 well microplate. Following this pre-incubation, 4MBr-5 cells are added to the mixture. The assay mixture in a total volume of  $100\ \mu\text{L}$ , containing antibody at concentrations of  $0.01$  to  $100\ \mu\text{g/mL}$ , recombinant human FGF-7 (KGF) at  $125\ \text{ng/mL}$ , and cells at  $1.5 \times 10^5$  cells/mL, is incubated at  $37^{\circ}\text{C}$  for 48 hours in a humidified  $\text{CO}_2$  incubator. The mixture is pulsed with  $^3\text{H}$ -thymidine during the final 24 hours. The cells are detached and harvested onto glass fiber filters, and the  $^3\text{H}$ -thymidine incorporated into the DNA is measured.

The Neutralization Dose<sub>50</sub> (ND<sub>50</sub>) for Anti-FGF-7 is  $\sim 6-12\ \mu\text{g/mL}$  in the presence of  $125\ \text{ng/mL}$  of recombinant human FGF-7 using the 4MBr-5 cell line.

The ND<sub>50</sub> is the concentration of antibody required to yield one-half maximal inhibition of the cytokine activity on a responsive cell line, when the cytokine is present at a concentration just high enough to elicit a maximum response.

The exact concentration of antibody required to neutralize human FGF-7 activity is dependent on the cytokine concentration, cell type, growth conditions, and the type of activity studied.

Immunoblotting: a working concentration of  $0.1-0.2\ \mu\text{g/mL}$  is recommended. The detection limit for recombinant human FGF-7 is  $\sim 1\ \text{ng/lane}$  and  $0.5\ \text{ng/lane}$  under non-reducing and reducing conditions, respectively.

ELISA: a working concentration of  $0.5-1.0\ \mu\text{g/mL}$  is recommended. The detection limit for recombinant human FGF-7 is  $\sim 0.08\ \text{ng/well}$ .

**Note:** In order to obtain the best results in various techniques and preparations, we recommend determining optimal working dilutions by titration.

Endotoxin level is  $< 0.1\ \text{EU}$  (endotoxin units) per  $1\ \mu\text{g}$  antibody as determined by the LAL (Limulus amoebocyte lysate) method.

### References

1. Givol, D., and Yayon, A., *FASEB. J.*, **6**, 3362 (1992).
2. Baird, A., et al., *Curr. Opin. Neurobiol.*, **4**, 78 (1994).
3. Fernig, F.G., and Gallagher, J.T., *Prog. Growth Factor Res.*, **5**, 353 (1994).
4. Kirkoshi, J., et al., *Biochem. Biophys. Res. Commun.*, **274**, 337 (2000).
5. Nishimura, T., et al., *Biochem. Biophys. Acta.*, **1492**, 203 (2000).
6. Bernard, O., and Matthew, P., *Guidebook to Cytokines and Their Receptors*, Oxford Press (New York, NY, 1994).
7. Galzie, Z., et al., *Biochem. Cell Biol.*, **75**, 669 (1997).
8. Callard, R., and Gearing, A., *The Cytokine Facts Book*, Academic Press (New York, NY, 1994).
9. Finch, P.W., et al., *Science*, **245**, 752 (1989).

KAA,PHC 01/08-1