



Product Information

Bid **Mouse, Recombinant**

Product Number **B8306**
Storage Temperature $-20\text{ }^{\circ}\text{C}$

Product Description

Mouse Bid (amino acids 1-195, accession number NM_007544) is expressed in *E. coli* and purified from the soluble fraction of disrupted cells. Purification did not employ detergents. Recombinant mouse Bid migrates as a polypeptide of 22 kDa on SDS reducing and non-reducing PAGE. A faint band corresponding to a disulfide-linked dimer is observed with non-reducing PAGE. Greater than 95% of the mouse Bid elutes as a 27 kDa protein on size exclusion chromatography, indicating that greater than 95% of the recombinant mouse Bid is monomeric.

Bid is a member of the Bcl-2 family that regulates outer mitochondrial membrane permeability.¹ Bid is a pro-apoptotic that can cause cytochrome c to be released from the mitochondrial intermembrane space into the cytosol. In healthy cells Bid is cytosolic. In response to Fas ligand or TNF, Bid is cleaved by caspase-8 and then relocates to the mitochondria outer membrane.^{2,3} Cleavage of Bid by caspase-8 generates a new N-terminal glycine, which apparently is myristoylated to target Bid to the mitochondria.⁴ Bid may then interact with another pro-apoptotic Bcl-2 family member, Bak.⁵ The interaction of Bid with Bak causes altered mitochondrial membrane permeability. Binding to the anti-apoptotic member Bcl-X_L neutralizes the activity of Bid.

The typical EC₅₀ for cytochrome c releasing activity is between 50 and 300 nM. The EC₅₀ for the desired application should be determined.

Purity: >95% (SDS-PAGE)

The product is supplied frozen as a 0.2 μm filtered solution in 25 mM HEPES, pH 7.5, containing 0.1 M KCl.

Precautions and Disclaimer

This product is for laboratory use only. Please consult the Material Data Safety Sheet for information regarding hazards and safe handling practices.

Storage/Stability

Stable for at least one year when store at $-20\text{ }^{\circ}\text{C}$. After thawing, sterile solutions may be stored at $4\text{ }^{\circ}\text{C}$ for one week or aliquoted under sterile conditions and stored at $-20\text{ }^{\circ}\text{C}$. Avoid repeated freeze-thaw cycles.

References

1. Gross, A. *et al.*, BCL-2 family members and the mitochondria in apoptosis. *Genes and Develop.*, **13**, 1899-1911 (1999).
2. Luo, X. *et al.*, Bid, a Bcl2 interacting protein, mediates cytochrome c release from mitochondria in response to activation of cell surface death receptors. *Cell*, **94**, 481-490 (1998).
3. Li, H. *et al.*, Cleavage of BID by caspase 8 mediates the mitochondrial damage in the Fas pathway of apoptosis. *Cell*, **94**, 491-501 (1998).
4. Zha, J. *et al.*, Posttranslational N-myristoylation of BID as a molecular switch for targeting mitochondria and apoptosis. *Science*, **290**, 1761-1765 (2000).
5. Wei, M.C. *et al.*, tBID, a membrane-targeted death ligand, oligomerizes BAK to release cytochrome c. *Genes Dev.*, **14**, 2060-2071 (2000).

RBR/AGW 12/01

Sigma brand products are sold through Sigma-Aldrich, Inc.

Sigma-Aldrich, Inc. warrants that its products conform to the information contained in this and other Sigma-Aldrich publications. Purchaser must determine the suitability of the product(s) for their particular use. Additional terms and conditions may apply. Please see reverse side of the invoice or packing slip.