



3050 Spruce Street
Saint Louis, Missouri 63103 USA
Telephone 800-325-5832 • (314) 771-5765
Fax (314) 286-7828
email: techserv@sial.com
sigma-aldrich.com

Product Information

Anti-CENP-E

Developed in Rabbit
IgG Fraction of Antiserum

Product Number **C 7488**

Product Description

Anti-CENP-E is developed in rabbit using as immunogen a synthetic peptide corresponding to amino acids 2545-2563 located near the C-terminus of human CENP-E conjugated to KLH. Whole antiserum is fractionated and then further purified by ion-exchange chromatography to provide the IgG fraction of antiserum that is essentially free of other rabbit serum proteins.

Anti-CENP-E recognizes human CENP-E (~310 kDa). Applications include the detection of CENP-E by immunoblotting and immunocytochemistry. Staining of CENP-E in immunoblotting is specifically inhibited with the CENP-E immunizing peptide (human, amino acids 2545-2563).

Chromosome movements during mitosis are orchestrated primarily by the interaction of mitotic spindle microtubules with the kinetochore, the site of attachment of spindle microtubules to the centromere.¹ Centromere-associated protein-E (CENP-E, 312 kDa)² is a member of the kinesin superfamily of microtubule motor proteins and is an integral part of kinetochore corona fibers that link centromere to the spindle microtubules.^{3,4} CENP-E plays an important role in attachment of kinetochores to spindle microtubules in the alignment of chromosomes and is an essential component of mitotic checkpoint signaling cascade.⁴⁻⁸ CENP-E localizes to the kinetochore throughout all phases of mitotic chromosome movement from early prometaphase through anaphase A. Cell-cycle dependent accumulation of CENP-E yields a maximum of 5,000 molecules per HeLa cell at the G2/M-phase transition.

Several studies indicate that CENP-E functions as a motor in the initial chromosome movement at the mitotic midzone. Depletion of CENP-E from *Xenopus* egg extracts disrupts metaphase chromosome alignment. CENP-E has been shown to be associated

with minus end-directed microtubule motor activity suggesting that CENP-E might be responsible for poleward kinetochore movements in prometaphase and anaphase A.⁵ In addition, CENP-E powers movement toward microtubule plus-end suggesting that it functions at the trailing kinetochore to link antipoleward movement to microtubule growth.⁶ Suppression of CENP-E synthesis by antisense CENP-E yields chromosomes that are chronically mono-oriented with flattened bipolar spindles and generates spindle poles fragments. Depletion of CENP-E leads to profound checkpoint activation and long-term mitotic arrest. CENP-E has been implicated as a binding partner for the mitotic checkpoint kinase BubR1 during mitosis.⁸⁻¹⁰

Reagent

Anti-CENP-E is supplied as a solution in 0.01 M phosphate buffered saline, pH 7.4, containing 15 mM sodium azide.

Precautions and Disclaimer

Due to the sodium azide content, a material safety data sheet (MSDS) for this product has been sent to the attention of the safety officer of your institution. Consult the MSDS for information regarding hazards and safe handling practices.

Storage/Stability

For continuous use, store at 2-8 °C for up to one month. For prolonged storage, freeze in working aliquots at -20 °C. Repeated freezing and thawing is not recommended. Storage in frost-free freezers is also not recommended. If slight turbidity occurs upon prolonged storage, clarify the solution by centrifugation before use. Working dilutions should be discarded if not used within 12 hours.

Product Profile

For immunoblotting, a minimum working antibody dilution of 1:2,000 is recommended using a whole cell extract of nocodazole-treated human epidermoid carcinoma A431 cell line.

For immunocytochemistry, a minimum working antibody dilution of 1:100 is recommended using the human epitheloid carcinoma HeLa cell line.

Note: In order to obtain the best results using various techniques and preparations, we recommend determining the optimal working dilutions by titration.

References

1. Scholey, J.M., et al., J. Cell Biol., **154**, 261-266 (2001).
2. Sharp, D.J., et al., Nature, **407**, 41-47 (2000).
3. Yen, T.J., et al., Nature, **359**, 536-539 (1992).
4. Yao, X., et al., J. Cell Biol., **139**, 435-446 (1997).
5. Thrower, D.A., et al., EMBO J., **14**, 918-926 (1995).
6. Wood, K.W., et al., Cell, **91**, 357-366 (1997).
7. Schaar, B.T., et al., J. Cell Biol., **139**, 1373-1382 (1997).
8. Yao, X., et al., Nature Cell Biol., **2**, 484-491 (2000).
9. Chan, G.K.T., et al., J. Cell Biol., **143**, 49-63 (1998).
10. Abrieu, A., et al., Cell, **102**, 817-826 (2000).

ER/KAA 07/03

Sigma brand products are sold through Sigma-Aldrich, Inc.

Sigma-Aldrich, Inc. warrants that its products conform to the information contained in this and other Sigma-Aldrich publications. Purchaser must determine the suitability of the product(s) for their particular use. Additional terms and conditions may apply. Please see reverse side of the invoice or packing slip.