



Product Information

ANTI-EPHRIN-A1, MOUSE, Developed In Goat, Affinity Isolated Antibody

Product Number **E7150**

Product Description

Anti-Ephrin-A1 is developed in goat using purified recombinant mouse ephrin-A1 extracellular domain, expressed in mouse NSO cells, as immunogen. The antibody is purified using mouse ephrin-A1 affinity chromatography.

Anti-Ephrin-A1 recognizes recombinant mouse ephrin-A1 in ELISA and immunoblotting. By ELISA and immunoblotting, the antibody shows approximately 50% cross-reactivity with recombinant human ephrin-A5, and 1% cross-reactivity with recombinant mouse ephrin-A2, recombinant human ephrin-A3, recombinant human ephrin-A4, recombinant mouse ephrin-B1, recombinant mouse ephrin-B2 and recombinant mouse ephrin-B3. By immunoblotting no cross-reactivity is shown with recombinant human EphA1, recombinant mouse EphA3, recombinant mouse EphA4 and recombinant mouse EphB6.

Anti-Ephrin-A1 may be used for ELISA and immunoblotting.

Eph receptor tyrosine kinases (RTK)¹ belong to a distinct class of RTKs that play important roles in many tissues. There are two classes of receptors, designated A and B. Both the A and B class receptors have an extracellular region consisting of a globular domain, a cysteine-rich domain, and two fibronectin type III domains. The transmembrane region and cytoplasmic region follow this. The cytoplasmic region contains a juxtamembrane motif with two tyrosine residues, which are the major autophosphorylation sites, a kinase domain, and a conserved sterile alpha motif (SAM) in the carboxy tail which contains one conserved tyrosine residue. Activation of kinase activity occurs after ligand recognition and binding. To date, at least 14 members of the Eph receptor family and a family of 8 ligands have been identified. Ligands of Eph family receptors are structurally related membrane-bound proteins that can be subdivided into two major subclasses,² ephrin-A

and ephrin-B. Ligands in the ephrin-A subclass, including the prototype family member ephrin-A1 (B61), are membrane associated through glycosylphosphatidylinositol linkages, whereas ephrin-B subclass consists of ligands with transmembrane domains. The a general role of the Eph family is in mediating repulsive cell-cell interaction, as suggested by studies of axonal guidance³⁻⁶ and neural crest cell migration.⁷⁻⁹

Reagents

Anti-Ephrin-A1 is supplied lyophilized from a 0.2 μm filtered solution of phosphate buffered saline. Endotoxin level is < 10 ng per mg antibody as determined by the LAL method.

Preparation Instructions

To one vial of lyophilized powder, add 1 ml of 0.2 μm filtered PBS to produce a 0.1 mg/ml stock solution of antibody. If aseptic technique is used, no further filtration should be needed for use in cell culture environments.

Storage/Stability

Prior to reconstitution, store at -20°C. Reconstituted product may be stored at 2-8°C for at least one month. For prolonged storage, freeze in working aliquots at -20°C. Avoid repeated freezing and thawing

Product Profile

For indirect ELISA, a working concentration of 0.5-1.0 μg/ml is determined to detect a limit of ~0.3 ng/well of recombinant mouse ephrin-A1.

For indirect immunoblotting, a working concentration of 0.1-0.2 μg/ml is determined using mouse ephrin-A1 at 5 ng/lane under non-reducing and reducing conditions.

Note: In order to obtain best results in different techniques and preparations we recommend determining optimal working dilutions by titration test.

References

1. Fraser, S., et al., Nature, **344**, 431 (1990).
2. Gale, N. W., et al., Neuron, **17**, 9 (1996).
3. Cheng, H., et al., Cell, **82**, 371 (1995).
4. Drescher, U., et al., Cell, **82**, 359 (1995).
5. Nakamoto, M., et al., Cell, **86**, 755 (1996).
6. Henkemeyer, M., et al., Cell, **86**, 35 (1996).
7. Krull, C. E., et al., Curr. Biol., **7**, 571 (1997).
8. Smith, A., et al., Curr. Biol. **7**, 561 (1997).
9. Wang, H. U., and Anderson, D. J., Neuron, **18**, 383 (1997).

jo/lpg 3/00

Sigma brand products are sold through Sigma-Aldrich, Inc.

Sigma-Aldrich, Inc. warrants that its products conform to the information contained in this and other Sigma-Aldrich publications. Purchaser must determine the suitability of the product(s) for their particular use. Additional terms and conditions may apply. Please see reverse side of the invoice or packing slip.