




Product Information

SIGMA QUALITY CONTROL TEST PROCEDURE

Enzymatic Assay of LUCIFERASE (EC 1.13.12.7) from Firefly

PRINCIPLE:



Abbreviations used:

ATP = Adenosine 5'-Triphosphate

AMP = Adenosine 5'-Monophosphate

CONDITIONS: T = 25°C, pH = 7.6

METHOD: Luminometric

REAGENTS:

- A. 1 M Glycine-Tris Buffer with 10 mM Ethylenediaminetetraacetic Acid and 100 mM Magnesium Sulfate, pH 7.0 at 25°C (Enz Dil)
(Prepare 50 ml in deionized water using Glycine, Free Base, Sigma Prod. No. G-7126, Ethylenediaminetetraacetic Acid, Disodium Salt, Dihydrate, Sigma Stock No. ED2SS, and Magnesium Sulfate, Heptahydrate, Sigma Prod. No. M-1880. Adjust to pH 7.0 at 25°C with solid Trizma Base, Sigma Prod. No. T-1503.)
- B. 0.15 mM Luciferin, 50 mM glycine, 1 mM Tris, 5 mM MgSO₄, 0.5 mM Ethylenediaminetetraacetic Acid, 0.1% (w/v) Bovine Serum Albumin, and 0.1% (w/v) Sodium Azide
(Prepare by reconstituting 1 vial of Firefly Diluent, Glycine Diluent with added Luciferin, Sigma Prod. No. F-3766 with 10 ml of deionized water. Store on ice.)
- C. 1 mM Adenosine 5'-Triphosphate Solution
(Prepare 10 ml in deionized water using Adenosine 5'-Triphosphate, Disodium Salt, Sigma Prod. No. A-5394.)
- D. 0.01 mM Adenosine 5'-Triphosphate Solution (ATP)
(Prepare 10.0 ml by adding 0.1 ml of Reagent C to 9.9 ml of deionized water.)

**Enzymatic Assay of LUCIFERASE
(EC 1.13.12.7)
from Firefly**

REAGENTS: (continued)

- E. 0.15 mM Luciferin, 50 mM Glycine, 1 mM Tris, 5 mM Magnesium Sulfate, 0.5 mM Ethylenediaminetetraacetic Acid, 0.1% (w/v) Bovine Serum Albumin, 0.1% (w/v) Sodium Azide, and 100 nM Adenosine 5'-Triphosphate Solution (Reaction Cocktail)
(Immediately before use, prepare by adding 0.010 ml of Reagent D to 0.99 ml of Reagent B.)
- F. ^{14}C 1.0 μCi Standard (^{14}C Std)
(Use ^{14}C 1.0 μCi Counting Standard.)¹
- G. Luciferase Enzyme Solution
(Immediately before use, prepare a solution containing 3,000,000 - 4,000,000 units/ml of Luciferase in cold Reagent A.)

PROCEDURE:

Set Reagent F (^{14}C Std) equal to 50% on the 100 scale of a photometer. Pipette (in milliliters) the following reagents into suitable containers:

	<u>Test</u>
Reagent E (Reaction Cocktail)	0.050

Incubate at 25°C for 10 minutes. Then add:

Reagent G (Enzyme Solution)	0.001
-----------------------------	-------

Immediately after adding Reagent G (Enzyme Solution), shake the tube twice and record the peak height of the luminescence.

CALCULATIONS:

$$\text{Units/ml enzyme} = \frac{(\text{Photometer Reading})(df)}{(0.001)}$$

Photometer Reading = One light unit will produce one count at the peak height of luminescence

df = Dilution factor

0.001 = Volume (in milliliter) of enzyme used

**Enzymatic Assay of LUCIFERASE
(EC 1.13.12.7)
from Firefly**

CALCULATIONS: (continued)

$$\text{Units/mg solid} = \frac{\text{units/ml enzyme}}{\text{mg solid/ml enzyme}}$$

$$\text{Units/mg protein} = \frac{\text{units/ml enzyme}}{\text{mg protein/ml enzyme}}$$

UNIT DEFINITION:

Light units are measured in 50 μl of assay mixture containing 5 picomoles of ATP and 7.5 nanomoles luciferin in glycine Tris buffer, pH 7.6 at 25°C. One light unit produces a biometer peak height equivalent to 0.02 μCi of ^{14}C in PPO/POPOP cocktail.

FINAL ASSAY CONCENTRATION:

In a 0.051 ml reaction mix, the final concentrations are 0.15 mM luciferin, 49 mM glycine, 0.98 mM Tris, 7 mM magnesium sulfate, 0.7 mM ethylenediaminetetraacetic acid, 0.1% (w/v) bovine serum albumin, 100 nM adenosine 5'-triphosphate and 3,000 - 4,000 units luciferase.

REFERENCE:

Leach, F.R. and Webster, J.J. (1986) *Methods in Enzymology*, 133, Part B, 51-70

Lin, S. and Cohen, H.P. (1968) *Analytical Biochemistry* **24**, 531-540

Strehler, B.L. (1974) in *Methods of Enzymatic Analysis* (Bergmeyer, H.U. ed.) 2nd ed., Vol. 4, 2112-2121

NOTES:

1. The ^{14}C 1.0 μCi Counting Standard used was obtained from Johnston's Lab Inc. (Prod. No. LS-1). The scintillation cocktail used is PPO(2,5-Diphenyloxazole)/POPOP(1,4-bis[5-Phenyl-2-oxazolyl]-benzene). This standard is no longer available. Other acceptable counting standards may be used.
2. This assay is based on the cited references.

**Enzymatic Assay of LUCIFERASE
(EC 1.13.12.7)
from Firefly**

NOTES: (continued)

3. Where Sigma Product or Stock numbers are specified, equivalent reagents may be substituted.

Sigma warrants that the above procedure information is currently utilized at Sigma and that Sigma products conform to the information in Sigma publications. Purchaser must determine the suitability of the information and products for its particular use. Upon purchase of Sigma products, see reverse side of invoice or packing slip for additional terms and conditions of sale.