



SIGMA-ALDRICH

3050 Spruce Street  
Saint Louis, Missouri 63103 USA  
Telephone 800-325-5832 • (314) 771-5765  
Fax (314) 286-7828  
email: techserv@sial.com  
sigma-aldrich.com

## Product Information

### Sudan Black B

Product Code **19,966-4**

Store at Room Temperature

Replacement for Product Number **S 0395**

#### Product Description

Molecular Formula:  $C_{29}H_{24}N_6$

Molecular Weight: 456.6

CAS Number: 4197-25-5

$\lambda_{max} = 596-605 \text{ nm}^1$

Extinction coefficient:  $E^{1\%} = 575-630$  (596-605 nm, ethanol)

Synonyms: Solvent Black B1, Fat Black HB, Solvent Black 3

Sudan Black B was found to eliminate lipofuscin-like autofluorescence in mammalian neural tissue without adversely affecting other fluorescent labels.<sup>2</sup> It was also found to reduce autofluorescence in archival formaldehyde-fixed paraffin-embedded myocardium tissue.<sup>3</sup>

Sudan Black B was used to detect native and oxidized low density lipoproteins (LDLs) after separation by capillary isotachopheresis (CITP).<sup>4</sup>

It has been used for staining fat in animal tissues and in bacteria (Burdon's method),<sup>5</sup> as well as the lipid portion of lipoprotein in polyacrylamide gel electrophoresis.<sup>6,7,8</sup>

#### Precautions and Disclaimer

For Laboratory Use Only. Not for drug, household or other uses.

#### Preparation Instructions

Sudan Black B is soluble in ethanol (10 mg/ml), yielding a dark blue solution. This may require heat for complete solubilization. It is also soluble in acetone, benzene, toluene, hydrocarbon solvents, fats, oils, and paraffins. It is slightly soluble in water (0.1 mg/ml).<sup>9</sup>

#### Procedure

To stain lipoprotein after polyacrylamide gel electrophoresis.<sup>7,8</sup>

1. Prepare a staining solution of 500 mg Sudan Black B in 20 ml of acetone. This is added to 15 ml of acetic acid, and then added to 85 ml of water.
2. Stir the mixture for 30 minutes and centrifuge to remove the precipitate.
3. Stain gel overnight in this solution.
4. Destain the gels in 3 changes of the following solution: 150 ml of acetic acid, 200 ml of acetone, 650 ml of water.

#### References

1. Conn's Biological Stains, Lillie, ed., p. 577 (1977).
2. Schnell S. A., et al., Reduction of lipofuscin-like autofluorescence in fluorescently labeled tissue. *J. Histochem. Cytochem.*, **47(6)**, 719-730 (1999).
3. Baschong, W. et al., Control of autofluorescence of archival formaldehyde-fixed, paraffin-embedded tissue in confocal laser scanning microscopy (CLSM). *J. Histochem. Cytochem.*, **49(12)**, 1565-1572 (2001).
4. Zorn, U. et al., Characterization of modified low density lipoprotein subfractions by capillary isotachopheresis. *Electrophoresis*, **22(6)**, 1143-1149 (2001).
5. Clark, G., *Staining Procedures*, 4th Ed., (Williams and Wilkins, Baltimore, 1981), pp. 193, 229, 393-394
6. Prat, J.P., et al., [Staining of lipoproteins after electrophoresis in polyacrylamide gel] *Bull. Soc. Chim. Biol.*, **51(9)**, 1367 (1969).
7. *Electrophoresis: Theory, Techniques, and Biochemical and Clinical Applications*, 2nd ed., ed. A.T. Andrews (Oxford University Press, Oxford, England, 1986), p. 37.
8. *Clinica Chimica Acta*, **203**, 109-188 (1991).
9. *The Sigma-Aldrich Handbook of Stains, Dyes & Indicators*, Green, F.J., Ed., Aldrich Chemical Co. (Milwaukee, WI: 1990), p. 660-661.

FEB/NSB 7/03

Sigma brand products are sold through Sigma-Aldrich, Inc.

Sigma-Aldrich, Inc. warrants that its products conform to the information contained in this and other Sigma-Aldrich publications. Purchaser must determine the suitability of the product(s) for their particular use. Additional terms and conditions may apply. Please see reverse side of the invoice or packing slip.