



SIGMA-ALDRICH

3050 Spruce Street
Saint Louis, Missouri 63103 USA
Telephone 800-325-5832 • (314) 771-5765
Fax (314) 286-7828
email: techserv@sial.com
sigma-aldrich.com

Product Information

TWEEN® 80 Sigma Ultra

Product Number **P 8074**
Store at Room Temperature

Product Description

CAS Number: 9005-65-6
Specific gravity: 1.07 (25 °C)
HLB (hydrophile-lipophile balance) value: 15.0^{1,2}
Critical Micellar Concentration (CMC): 13-15 mg/liter^{1,3}
Brookfield Viscosity: 400-620 cps (25 °C, neat)
Micelle molecular weight: 76 kDa³
Synonyms: Polysorbate 80, PEG (80) sorbitan monooleate, polyoxyethylenesorbitan monooleate

TWEEN 80 is a polyethylene sorbitol ester, with a calculated molecular weight of 1,310 daltons, assuming 20 ethylene oxide units, 1 sorbitol, and 1 oleic acid as the primary fatty acid.⁴ Fatty acid constituents of this product are determined by transesterification to yield fatty acid methyl esters, which are identified by gas chromatography. Typically the fatty acid composition is approximately 70% oleic acid with several other fatty acids such as palmitic acid indicated.

TWEEN 80 has been widely used in biochemical applications including: solubilizing proteins, isolating nuclei from cells in culture,⁵ growing of tubercule bacilli,⁶ and emulsifying and dispersing substances in medicinal and food products. It has little or no activity as an anti-bacterial agent¹ except it has been shown to have an adverse effect on the antibacterial effect of methyl paraben and related compounds.⁷ Polysorbates have been reported to be incompatible with alkalis, heavy metal salts, phenols, and tannic acid. They may reduce the activity of many preservatives.⁸

Product No. P 8074 is for applications which require tight control of elemental content. Trace elemental analyses have been performed on this SigmaUltra TWEEN 80. The Certificate of Analysis provides lot-specific results.

Sigma also offers these additional TWEEN 80 products.

P 1754 - for general use.

P 4675 and P 4780 - tested in cell culture applications.

P 5188 - tested in molecular biology applications.

P 6224 - non animal source.

Precautions and Disclaimer

For Laboratory Use Only. Not for drug, household or other uses.

Preparation Instructions

TWEEN 80 is miscible in water (0.1 ml/ml) yielding a clear to slightly hazy faint yellow solution. It is reported to be miscible with alcohol, cottonseed oil, corn oil, ethyl acetate, methanol, and toluene, but insoluble in mineral oil.⁴ The pH of a 1% aqueous solution is 5.5-7.2.

Storage/Stability

Aqueous solutions of polysorbates as well as the neat liquid will undergo autooxidation over time, with changes being catalyzed by light, increased temperature, and copper sulfate.⁹ Solutions are reasonably stable at 2 - 8 °C for short periods. For special applications, storage under argon or nitrogen may be preferred.

The product is not sterile. Autoclaving of solutions is generally not advised. Sterile filtration is more easily done if the liquid is warmed to about 40 °C and alternate portions of hot distilled water and TWEEN 80 are poured through the 0.22 µm filter. The TWEEN 80 will blend and remain in solution.

References

1. Data for Biochemical Research, 3rd ed., Dawson, R. M. C., et al., Oxford University Press (New York, NY: 1986), p. 289.
2. Neugebauer., J. M., Detergents: An overview. *Methods Enzymol.*, **182**, 247 (1990).
3. Protein Purification Methods: A Practical Approach., Harris, E. L. V., and Angal, S., eds., IRL Press at Oxford University Press (New York, NY: 1990), p. 71.
4. The Merck Index, 11th Ed., Entry #7559.
5. Fisher, H. W. and H. Harris, *Proc. R. Soc. B*, **156**, 521 (1962).
6. Dubos, R. J. and B.D. Davis, *J. Exp. Med.*, **83**, 409 (1946).

7. Disinfection, Sterilization & Preservation, 4th Ed., Block, S.S. (Lea & Febiger Pub., 1991) Chapter 4.
8. Martindale: The Extra Pharmacopoeia, 30th Ed., Reynolds, J. E. F., ed., Pharmaceutical Press(London, England: 1993) p. 1030.

9. Donbrow, M., et al., Autoxidation of polysorbates. J. Pharm. Sci., **67**, 1676-1681 (1978).

TWEEN is a registered trademark of Uniqema, a business unit of ICI Americas Inc.

RLG 5/06

Sigma brand products are sold through Sigma-Aldrich, Inc.

Sigma-Aldrich, Inc. warrants that its products conform to the information contained in this and other Sigma-Aldrich publications. Purchaser must determine the suitability of the product(s) for their particular use. Additional terms and conditions may apply. Please see reverse side of the invoice or packing slip.