



Product Information

MONOCLONAL ANTI-PSTAIR

Clone PSTAIR

Mouse Ascites Fluid

Product Number **P 7962**

Product Description

Monoclonal Anti-PSTAIR (mouse IgG1 isotype) is derived from the PSTAIR hybridoma produced by the fusion of mouse myeloma cells and splenocytes from BALB/c mice. A synthetic 16 amino acid oligopeptide containing the PSTAIR sequence, conjugated to BSA, was used as the immunogen.¹ The isotype is determined using Sigma ImmunoType™ Kit (Product Code ISO-1) and by a double diffusion immunoassay using Mouse Monoclonal Antibody Isotyping Reagents (Product Code ISO-2).

Cell division is a fundamental biological process, consisting of the splitting of the cell and its genetic material into two daughter cells. Mitosis results in the formation of two new nuclei, each having the same number of chromosomes as the parental nucleus. During the cell cycle of most somatic cells, DNA synthesis (S-phase) and mitosis (M-phase) are separated by two "growth" stages (G₁ and G₂) of varying duration. A typical eukaryotic cell sequentially passes through G₁, S, G₂, and M and back into G₁ during a single cycle.⁹ Maturation-promoting factor (MPF), originally found during meiosis in frog oocytes, is a cytoplasmic factor, which is highly conserved among a wide range of species and plays a key role in the progression of the cell cycle from interphase (G₂) to metaphase (M), in both meiosis and mitosis. The cell cycle can be considered as a cyclin-dependent kinases (cdks) cycle, which is probably controlled by biochemical modifications such as phosphorylation of CDKs and formation of complex(es) with other proteins, including the cyclins.¹⁰ CDKs are key regulators of cell cycle progression.^{11,12} CDKs are closely related in size (35-40 kDa) and sequence (>40% identity) and associate with and are activated by a cyclin, which acts as a regulatory subunit. In the fission yeast *Shizosaccharomyces pombe*, a single major CDK has been identified (cdc2) while in human cells, the growing list of CDKs now includes cdc2 (Cdk1), and Cdk2 to Cdk7.

The typical CDK catalytic subunit contains a 300 amino acid catalytic core that is completely inactive when monomeric and unphosphorylated.¹³ The primary regulator of CDK activity is the cyclin molecule. Each CDK interacts with a specific subset of cyclins which activate them by enabling their phosphorylation at specific residues. For example, the activation of cdc2 requires the phosphorylation of a conserved threonine 161 (Thr 160 in cdk2) and then it may be inactivated by a second phosphorylation at Tyrosine 15.¹⁴ CDKs are constitutively expressed throughout the cell cycle and are activated and inactivated by specific kinases and phosphatases which are responsible for the specific phosphorylation and dephosphorylation.^{15,16} In every eukaryote examined, CDKs contain an evolutionary conserved 16 amino acid sequence called PSTAIR (EGVPSTAIRESLLKE) which distinguishes them from other protein kinases. The PSTAIR motif is involved in the complex formation with cyclins. The availability of monoclonal antibody reacting specifically with the PSTAIR sequence^{1,17} enables the subcellular detection and localization of the various CDKs and examination of substrate interactions, in a variety of organisms.

Reagents

The product is provided as ascites fluid with 0.1% sodium azide as a preservative.

Precautions and Disclaimer

Due to the sodium azide content a material safety sheet (MSDS) for this product has been sent to the attention of the safety officer of your institution. Consult the MSDS for information regarding hazards and safe handling practices.

Storage/Stability

For extended storage freeze in working aliquots. For continuous use, store at 2-8 °C for up to one month. Repeated freezing and thawing is not recommended. Storage in "frost-free" freezers is not recommended. If slight turbidity occurs upon prolonged storage, clarify the solution by centrifugation before use.

Product Profile

Monoclonal Anti-PSTAIR recognizes an epitope present in the PSTAIR sequence¹ of p34^{cdc2} (cdk1) and of other cyclin-dependent kinases containing the PSTAIR motif (cdk2 and cdk3). The antibody recognizes 31-34 kDa proteins (1-4 bands) in immunoblotting.¹⁻⁷ The product is also reactive in ELISA¹ and immunoprecipitation.¹ It cannot precipitate p34^{cdc2} when it is complexed with cyclin B.¹ Cross-reactivity has been observed with human, monkey, mouse,^{1,8} Japanese quail,^{1,3} fish (goldfish,⁴⁻⁷ carp,^{1,5,7} eel,¹ starfish,¹ and amago salmon¹), *Xenopus*,¹ ciliates,¹ and plants (lily,^{1,2} and onion¹).

Monoclonal Anti-PSTAIR may be used for the localization of cyclin-dependent kinases, containing the PSTAIR motif using various immunochemical assays including ELISA, immunoblot and immunoprecipitation.

In order to obtain best results, it is recommended that each user determine the optimal working dilution for individual applications by titration assay.

References

1. Yamashita, M., et al., *Dev. Growth Differ.*, **33**, 617 (1991).
2. Yamaguchi, A., et al., *Dev. Growth Differ.*, **33**, 625 (1991).
3. Mori, M., et al., *Dev. Biol.*, **146**, 246 (1991).
4. Hirai, T., et al., *Dev. Biol.*, **152**, 113 (1992).
5. Yamashita, M., et al., *Eur. J. Biochem.*, **205**, 537 (1992).
6. Hirai, T., et al., *Mol. Reprod. Dev.*, **33**, 131 (1992).
7. Yamashita, M., et al., *Dev. Biol.*, **149**, 8 (1992).
8. Yasuda, H., et al., *Somat. Cell Mol. Genet.*, **18**, 403 (1992).
9. Freeman, R., and Donoghue, D., *Biochemistry*, **30**, 2293 (1991).
10. Nurburg, C., and Nurse, P., *Ann. Rev. Biochem.*, **61**, 441 (1992).
11. Heichman, K., and Roberts, J., *Cell*, **79**, 557 (1994).
12. Hunter, T., and Pines, J., *Cell*, **79**, 573 (1994).
13. Draetta, G., *Trends Cell Biol.*, **3**, 287 (1993).
14. Morgan, D., *Nature*, **374**, 131 (1995).
15. Mueller, P., et al., *Science*, **270**, 86 (1995).
16. Poon, R., and Hunter, T., *Science*, **270**, 90 (1995).
17. Kamo, K., et al., *J. Immunol. Meth.*, **156**, 163 (1992).

KMR/MAM 01/02

Sigma brand products are sold through Sigma-Aldrich, Inc.

Sigma-Aldrich, Inc. warrants that its products conform to the information contained in this and other Sigma-Aldrich publications. Purchaser must determine the suitability of the product(s) for their particular use. Additional terms and conditions may apply. Please see reverse side of the invoice or packing slip.