

Product Information

Hexokinase from *Saccharomyces cerevisiae*

Catalog Number **H5000**
 Storage Temperature $-20\text{ }^{\circ}\text{C}$

CAS RN 9001-51-8
 EC 2.7.1.1
 Synonyms: ATP:D-hexose 6-phosphotransferase

Product Description

Glycolysis is the process occurring in almost all living organisms by which they metabolize D-glucose to generate energy and metabolic intermediates. In the first step of glycolysis, hexokinase phosphorylates the C6 position of D-glucose in the presence of ATP by the following reaction:



Yeast hexokinase has three isozymes, designated P-I, P-II, and glucokinase (Glk1). Each has distinctive properties.¹ Yeast hexokinase P-II has both a catalytic and a regulatory function.²

pI:³

P-I	5.25
P-II	4.93

Several other hexoses can serve as substrates for hexokinase (relative reaction rates):⁴

glucose	1.0
2-deoxy-2-fluoro-D-glucose	0.5
mannosamine	0.2
5-thioglucoase	0.01
3-deoxy-3-aminoglucoase	0.003

Also:^{1,4} 1,5-anhydro-D-glucitol, 1-thio-D-glucose

K_M (mM):^{1,4}

D-glucose	0.12 (P-I and P-II)
D-fructose	0.33 (P-I and P-II)
D-mannose	0.04 (Glk1)
D-mannosamine	5
5-thio-D-glucose	4

Molecular Weight:⁵ 110 kDa (dimer)

Hexokinase is a dimeric protein with two equal 55 kDa monomers.

Extinction coefficient:

P-I	$E_{280}^{1\%} = 8.85$
P-II	$E_{280}^{1\%} = 9.47$

pH Optimum:⁶ 7.5–9.0

Activators: Mg^{2+} ($K_M = 2.6\text{ mM}$),
 catecholamine-related compounds⁷

Inhibitors ^{1,4}	K_i (mM)
D-glucosamine	1.5
D-mannose	0.06
D-xylose	25 (isozyme P-I) 80 (isozyme P-II)
6-deoxy-D-glucose	50
N-acetylmannosamine	50

Also sorbose-1-phosphate, polyphosphates, 6-deoxy-6-fluoroglucoase, 2-C-hydroxy-methylglucose, lyxose, and thiol reactive compounds⁶

Hexokinase is used for the determination of D-glucose, D-fructose, and D-sorbitol in food or other biological materials. Hexokinase can also be used in the assay of glycosides that are convertible to glucose or fructose.

This product (Catalog Number H5000) is purified from *Saccharomyces cerevisiae* (baker's yeast). It is supplied as a lyophilized powder. It is a mixture of isozymes.

Protein: ≥85%, balance primarily sodium citrate.

Specific activity: ≥25 units/mg protein (biuret)

Unit Definition: One unit will phosphorylate 1.0 μmole of D-glucose per min at pH 7.6 at 25 °C.

Hexokinase is assayed spectrophotometrically in a 2.57 ml reaction mixture containing 39 mM triethanolamine, 216 mM D-glucose, 0.74 mM ATP, 7.8 mM MgCl₂, 1.1 mM β-NADP, 2.5 units glucose-6-phosphate dehydrogenase, and 0.025–0.05 unit hexokinase.

Contaminants:

glucose-6-phosphate dehydrogenase (≤10%)

phosphoglucose isomerase (≤10%)

Precautions and Disclaimer

This product is for R&D use only, not for drug, household, or other uses. Please consult the Material Safety Data Sheet for information regarding hazards and safe handling practices.

Preparation Instructions

Hexokinase is soluble in cold water (0.5-1.0 unit/ml) or citrate buffer, pH 7.0.

Storage/Stability

Store the product at –20 °C and it is stable for at least 2 years when stored properly.

Solutions in water or citrate buffer have remained stable during repeated freezing and thawing for a period of 30 days.

References

1. Fernandez, R., *et al.*, Inhibition and inactivation of glucose-phosphorylating enzymes from *Saccharomyces cerevisiae* by D-xylose. *J. Gen. Microbiol.*, **131**, 2705 (1985).
2. Entian, K.D., and Fröhlich, K.U., *Saccharomyces cerevisiae* mutants provide evidence of hexokinase P-II as a bifunctional enzyme with catalytic and regulatory domains for triggering carbon catabolite repression. *J. Bacteriol.*, **158**, 29-35 (1984).
3. Schmidt, J.J., and Colowick, S.P., Identification of a peptide sequence involved in association of subunits of yeast hexokinases. *Arch. Biochem. Biophys.*, **158**, 471-77 (1973).
4. Machado de Domenech, E. E., and Sols, A., Specificity of hexokinases towards some uncommon substrates and inhibitors. *FEBS Lett.*, **119**, 174-176 (1980).
5. Jacob, L., *et al.*, Purification and crystallization of yeast hexokinase isozymes. Characterization of different forms by chromatofocusing. *J. Chromatogr.*, **587**, 85-92 (1991).
6. Sols, A., *et al.*, Substrate specificity and some other properties of baker's yeast hexokinase. *Biochem. Biophys. Acta*, **30**, 92-101 (1958).
7. Harrison, W.H., *et al.*, Aluminium inhibition of hexokinase. *Lancet*, **2**, 277 (1972).
8. Mulcahy, P., *et al.*, Application of kinetic-based biospecific affinity chromatographic systems to ATP-dependent enzymes: Studies with yeast. *Anal. Biochem.*, **309**, 279-92, (2002).

MB,JWM,MAM 12/06-1

Sigma brand products are sold through Sigma-Aldrich, Inc.

Sigma-Aldrich, Inc. warrants that its products conform to the information contained in this and other Sigma-Aldrich publications. Purchaser must determine the suitability of the product(s) for their particular use. Additional terms and conditions may apply. Please see reverse side of the invoice or packing slip.