

Product Information

Monoclonal Anti-Interleukin-4-Carboxyfluorescein Clone 3007.11

antibody produced in mouse, purified immunoglobulin

Catalog Number **F3901**

Product Description

Monoclonal Anti-Interleukin-4 (IL-4) (mouse IgG1 isotype) is derived from the hybridoma produced by the fusion of a mouse myeloma cell line and splenocytes from a mouse immunized with recombinant human IL-4. The conjugate is prepared by conjugation of carboxyfluorescein to Protein G purified IL-4 monoclonal antibody.

Monoclonal Anti-Interleukin-4-Carboxyfluorescein recognizes human interleukin-4 by flow cytometry.

Interleukin-4 (IL-4) is a lymphokine that has profound effects on the growth and differentiation of immunologically competent cells.¹ Synonyms for IL-4 include: B cell stimulatory factor-1 (BSF-1), T cell growth factor-2 (TCGF-2) and mast cell growth factor-2 (MCGF-2).²⁻⁴ Interleukin-4 is a complex glycoprotein released by a subset of activated T cells. The molecular weight of IL-4 occurring naturally is 12-20 kDa.

This product is designed to detect intracytoplasmic cytokines and enumerate cytokine-producing cells in a mixed population using flow cytometric assays. The cytokine profile of activated cells has profound implications on the immune status of the host. T helper cells are generally considered as the cellular sources of cytokines. In the original differentiation model for CD4+ T helper cells, precursors were referred to as Th₀ which differentiate into functional subsets designated as Th₁ and Th₂. Th₀ cells secrete a combination of both type I and type II cytokines, whereas Th₁ were thought to secrete only type I (IL-2, IL-12, TNF-β, and IFN-γ) cytokines and Th₂ only type II (IL-4, IL-5, IL-6, IL-10, IL-13) cytokines. In recent years additional experimental data have led to the revision of this model. The cytokine profiles of CD4+ T helper cells are not mutually exclusive as originally envisioned. In addition to CD4+ cells, CD8+ cells are also active in both type I and type II cytokine production. Clearly, the cytokine profile of specific functional cell subsets during immune activation influences the outcome of the response. The ability to manipulate the cytokine

production profile holds promise for modulating the balance between Th₁ and Th₂ or type I and type II cytokine effects towards achieving protective immunity. Such therapeutic potential, as well as the insight into underlying immune mechanisms, has stimulated much research into this area. Intracellular cytokine staining reagents can be used in flow cytometry to correlate cytokine production profiles at the single cell level with cell surface phenotype, e.g., CD4, CD8, activation and antigen expression. This approach is much more informative than the traditional methods of quantitation of secreted cytokine accumulated in culture supernatant of functionally heterogeneous cell populations. This method is rapid and avoids imprecise and tedious physical cell enrichment methods. Intracellular cytokine staining can be coupled with the Cell Census Plus™ System, Catalog Number CCPS1, to correlate the proliferation activity of immune cell subsets with particular cytokine-producing-profile(s) in response to different cell activation stimuli.

Reagent

Supplied as a solution in saline containing up to 0.5% bovine serum albumin and up to 0.1% sodium azide as preservative.

Precautions and Disclaimer

This product is for R&D use only, not for drug, household, or other uses. Please consult the Material Safety Data Sheet for information regarding hazards and safe handling practices.

Storage/Stability

Store at 2-8 °C. Protect from prolonged exposure to light. Do Not Freeze. If slight turbidity occurs upon prolonged storage, clarify the solution by centrifugation before use.

Procedure

Paraformaldehyde Fixative:

4.0 grams paraformaldehyde, Catalog Number P6148, in 100 mL sterile PBS, pH 7.4. Dissolve by heating to 56 °C for approx. 1 hour. Store 2-8 °C for up to 2 weeks. Protect from light.

Saponin Buffer:

0.1 gram saponin, Catalog Number S4521, dissolved in 100 mL of 1X Hank's Balanced Salt Solution (10X HBSS, Catalog Number H4641) with 0.05% sodium azide. Store at 22 °C for up to one month.

Conditions for Intracellular Cytokine Staining⁵⁻⁹

1. Cell preparation
Enrich peripheral blood for lymphocytes using ficoll. Stimulate lymphocytes *in vitro* with 10 ng/ml PMA (phorbol myristate acetate) and 1 μM calcium ionomycin. To inhibit protein secretion, add 2 μM monensin and culture for 18 hours. Harvest cells and wash in PBS.
2. Cell surface staining
If cells are to be stained with another monoclonal antibody conjugate, follow the manufacturer's staining procedure.
3. Fixation and permeabilization
Fix with cold 4% paraformaldehyde at 22 °C for 10 minutes, with intermittent vortexing to maintain a single cell suspension. Wash again and permeabilize with 0.1% saponin in buffer.
4. Antibody staining
To 1-5 x 10⁵ cells suspended in approximately 200 μL saponin buffer, add 10 μL Anti-IL-4-Carboxyfluorescein. Protect from light and incubate at 22 °C for 30-45 minutes. Wash cells 2 x with saponin buffer. Finally, resuspend cells in PBS containing 1% bovine serum albumin.
5. Detection
Analyze in a flow cytometer according to manufacturer's instructions.

Quality Control

It is advisable to run the appropriate negative controls. Negative controls establish background fluorescence and non-specific binding. The ideal negative control reagent is a mouse monoclonal or myeloma protein which has no reactivity with human cells. It should be isotype-matched to the antibody and of the same concentration and F/P molar ratio as the antibody. The degree of autofluorescence or negative control reagent fluorescence will vary with the type of cells under study and the sensitivity of the instrument used.

Product Profile

When assayed by flow cytometric analysis, using 10 μL of the antibody conjugate to stain 1-5 x 10⁵ cells, the antibody conjugate detects intracellular levels of cytokine in secreting cells.

Note: In order to obtain best results in different preparations, it is recommended that each individual user determine their optimum working dilutions by titration assay.

Specificity

The product is determined to be specific by a group of assays that include:

- a. Inhibition of staining in the presence of excess exogenously added IL-4.
- b. Inhibition of staining in the presence of unconjugated anti-IL-4 and/or
- c. Lack of staining when cells are fixed but not permeabilized

References

1. Howard, M., et al., *J. Exp. Med.*, **155**, 914 (1982).
2. Mosmann, T., et al., *Proc. Nat. Acad. Sci. USA*, **83**, 5654 (1986).
3. Paul, W., and Ohara, J., *Ann. Rev. Immunol.*, **5**, 429 (1987).
4. Park, L., et al., *J. Exp. Med.*, **166**, 476 (1987).
5. Jung, T., et al., *J. Immunol. Methods*, **159**, 197 (1993).
6. Prussin, C., and Metcalfe, D.D., *J. Immunol. Methods*, **188**, 117 (1995).
7. Prussin, C., *J. Clin. Immunol.*, **17**, 195 (1997).
8. Estcourt, C., *Clin. Immunol. Immunopathol.*, **83**, 60 (1997).
9. Ito, M., et al., *Clin. Immunol. Immunopathol.*, **83**, 281 (1997).

TD,KAA,PHC 07/09-2