

Product Information

Anti-Podoplanin (KS-17)

Developed in Rabbit
Affinity Isolated Antibody

Product Number **P 2120**

Product Description

Anti-Podoplanin (KS-17) is developed in rabbit using as immunogen a synthetic peptide corresponding to amino acid residues 65-81 of rat podoplanin, conjugated to KLH. The antibody is affinity-purified using the immunizing peptide immobilized on agarose.

Anti-Podoplanin (KS-17) recognizes rat podoplanin. Applications include immunoblotting (~38 kDa) and immunofluorescence staining. Detection of podoplanin by immunoblotting is specifically inhibited with the immunizing peptide.

Podoplanin is an integral membrane mucoprotein present on the surface of podocytes in kidney glomeruli and the parietal cells of the glomerular Bowman's capsule.¹⁻³ It is a single membrane-spanning protein composed of heavily glycosylated extracellular domain and a short intracytoplasmic tail. Glycoproteins with similar sequences were described in rodent lung type I alveolar cells, mesothelial cells, osteoblasts and osteocytes, and thymic epithelial cells.^{3,5} Podoplanin, also named E11 antigen, has been described to be present in normal, inflammatory, and neoplastic lymphatic capillary endothelium.^{1,3,4,6-9} It is not expressed *in vivo* in blood capillary endothelial cells, human hemangiomas, and lymph nodes high endothelial venules. Large lymphatic vessels having smooth muscle cells may lack podoplanin. Podoplanin is expressed selectively by dermal microvascular lymphatic endothelial cells but not by blood vascular endothelial cells. Expression of podoplanin is regulated by the lymphatic-specific homeobox gene Prox-1.⁷⁻⁹ Podoplanin does not seem to be upregulated in tumor angiogenesis. It is detectable in human angiosarcomas, lymphangiomas, and in Kaposi's sarcomas. Podoplanin appears to play an important role in the maintenance of podocyte foot processes, and hence, glomerular permeability.^{1,2} Experiments involving depletion of T1 α /podoplanin in mice indicate that it has a critical regulatory role in the formation of the lymphatic vasculature.⁷⁻⁹

Reagent

Anti-Podoplanin (KS-17) is supplied as a solution in 0.01 M phosphate buffered saline pH 7.4 containing 15 mM sodium azide.

Antibody Concentration: 1.2-1.5 mg/ml

Precautions and Disclaimer

Due to the sodium azide content, a material safety data sheet (MSDS) for this product has been sent to the attention of the safety officer of your institution. Consult the MSDS for information regarding hazards and safe handling practices.

Storage/Stability

For continuous use, store at 2-8 °C for up to one month. For prolonged storage, freeze in working aliquots at -20 °C. Repeated freezing and thawing is not recommended. Storage in frost-free freezers is also not recommended. If slight turbidity occurs upon prolonged storage, clarify the solution by centrifugation before use. Working dilutions should be discarded if not used within 12 hours.

Product Profile

By immunoblotting, a working antibody concentration of 2.5-5 μ g/ml is recommended using extracts of rat kidney glomeruli and rat lung and a chemiluminescent detection reagent.

By indirect immunofluorescence, a working antibody concentration of 10-20 μ g/ml is recommended to detect podoplanin by staining glomeruli in acetone-fixed rat kidney frozen sections.

Note: In order to obtain the best results using various techniques and preparations, we recommend determining the optimal working dilutions by titration.

References

1. Breiteneder-Geleff, S., et al., *Am. J. Pathol.*, **151**, 1141-1152 (1997).
2. Matsui, K., et al., *J. Am. Soc. Nephrol.*, **9**, 2013-2026 (1998).
3. Wetterwald, A, et al., *Bone*, **18**, 125-132 (1996).

4. Breiteneder-Geleff, S., et al., Am. J. Pathol., **154**, 385-394 (1999).
5. Zimmer, G., et al., Biochem. J., **341**, 277-284 (1999).
6. Kriehuber, E., et al., J. Exp. Med., **194**, 797-808 (2001).
7. Hong, Y.K., et al., Dev. Dyn., **225**, 351-357 (2002).
8. Schacht, V., et al., EMBO J., **22**, 3546-3556 (2003).
9. Petrova, T.V., et al., EMBO J., **21**, 4593-4599 (2002).

CAA/RM 02/04

Sigma brand products are sold through Sigma-Aldrich, Inc.

Sigma-Aldrich, Inc. warrants that its products conform to the information contained in this and other Sigma-Aldrich publications. Purchaser must determine the suitability of the product(s) for their particular use. Additional terms and conditions may apply. Please see reverse side of the invoice or packing slip.